

45

The effect of the presence of a freely diffusing compartment on observation of restricted diffusion in single- and double-PFG experiments

N. Shemesh¹, E. Özarslan², A. Bar-Shir¹, P. J. Basser², Y. Cohen¹;

¹Chemistry, Tel Aviv University, Tel Aviv, Israel, ²Section on Tissue Biophysics and Biomimetics, NICHD, National Institute of Health, Bethesda, MD, United States.

Introduction: Studies of restricted diffusion are commonly conducted using single pulsed-field-gradients (s-PFG) diffusion experiments¹. In homogeneous samples, the diffusion-diffraction phenomenon arising from a single population of diffusing species has been observed, and the restricted diffusion profile was used to extract important structural features from the sample². However, systems that are more realistic such as biological tissue and porous media are characterized by compartmentation which may complicate the interpretation of structural features. Double-PFG (d-PFG) experiments have lately been gaining interest due to their ability to extract small compartmental dimensions even at low q values³. Therefore, in this study, we characterized the superposition of restricted and free diffusion in s- and d-PFG both experimentally and theoretically using a novel composite bi-compartmental phantom, in which the „ground-truth“ is known *a-priori*.

Subjects and Methods: Figure 1 shows a cartoon of the bi-compartmental phantom. Freely diffusing water in the Fast-Diffusion-Compartment (FDC) undergoes Gaussian (free) diffusion while water in the microcapillaries experiences restricted diffusion forming the Slow-Diffusion-Compartment (SDC) of the bi-compartmental phantom.

Results: Figure 2 shows the s-PFG experiment conducted with varying diffusion periods on the bi-compartmental model. Two phenomena can be seen in the different q -regimes. For high q -values, the diffraction patterns from water diffusing in the SDC can be gradually observed. At $\Delta > 100$ ms, the diffusion profile doesn't change at high q -values. However, at lower q -values, e.g., $q < 200 \text{ cm}^{-1}$, the diffusion curves change dramatically for each value of Δ , and shows increased attenuation of the signal with increasing Δ . Solid lines represent the theoretical curves.

Figures 3A and 3B show the angular d-PFG experiment in the bi-compartmental model at low and high q -values respectively. At the low- q regime, the angular dependence is lost, and the size of the compartment cannot be accurately extracted. However, at higher q -values, the angular dependence is retained, and the accurate size of the compartment is extracted.

Conclusions: The effect of adding an FDC on the observation of restricted diffusion was studied using s-PFG and d-PFG experiments. Importantly, structural information can be extracted at higher q -values where the restricted diffusion is accentuated, and the free diffusion is suppressed. This may be of importance in biological tissue and porous media.

References:

1. E.O. Stejskal and J. E. Tanner, J. Chem. Phys. 42 (1965) 288-292.
2. P.T. Callaghan et al., Nature 351 (1991) 467-469
3. Shemesh et al., J. Magn. Reson. 198 (2008) 15-23

